

Chapter 2

Evolutionary Processes in Crop Species Enhanced by Agritech

Wisdom Leaf Press

Pages number, 6–10

© The Author 2024

<https://journals.icapsr.com/index.php/wlp>

DOI: 10.55938/wlp.v1i2.104



Mansi Sahu¹, Kailash Bisht² and Sanjeev Kumar Shah³

Abstract

Traditional breeding techniques, ethno-botanical expertise, local agronomy research, extension services, farmer engagement, and social and cultural studies are all still important, even if gene-focused domestication initiatives might produce vital allele variations for novel crop production. The scientific community, funding organizations, proposal reviewers, and researchers themselves must acknowledge and encourage these fields as essential to the advancement of gene editing technology in order to fully realize the benefits of de novo domestication on the environment and society. Classical genetic mapping has identified numerous genes and loci related to domestication and alterations. However, only a small percentage of these genes have been fully described. The genetic valley preserves advantageous haplotypes, resulting in less genetic variety. It is anticipated that advances in genetic mapping, crop genome sequencing, and various biological data collecting will strengthen our comprehension of crop biological processes and accelerate the conversion of lab results into practical field applications. A species' ability to adapt to external challenges is made possible by genetic variety. Since it permits modifications to the genetic makeup, animals can adjust to shifts in their surroundings. Enhancing morphological and agronomic traits is a major function of plant genetic diversity in agriculture. A greater degree of diversity improves a species' ability to adapt to changing circumstances, particularly when pests and climatic swings arise. With an emphasis on genetic variables and phenotypic plasticity, this study investigates diversity in seed dispersion attributes. It investigates certain characteristics and biological systems and suggests a simulation model to meet the demands of upcoming studies. The researchers call on biologists and ecologists to comprehend how fast changes in seed distribution affect plant population responses to climate change.

Keywords

Functional Genomics, Genetic Modification, Green Revolution, Non-Thermal Plasma, Phenotyping Technological Advances, Plant Molecular Breeding

¹Division of Research & Innovation, Uttarakhand University, Dehradun. mansi.smile.1999@gmail.com

²Uttarakhand Institute of Management, Uttarakhand University, Dehradun, India. kailash.bisht1911@gmail.com

³Uttarakhand Institute of technology, Uttarakhand University, Dehradun, Uttarakhand, India. sanjeevshah19@gmail.com

Corresponding author:

Email id: sanjeevshah19@gmail.com



© 2024 by Mansi Sahu, Kailash Bisht and Sanjeev Kumar Shah Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license, (<http://creativecommons.org/licenses/by/4.0/>). This work is licensed under a Creative Commons Attribution 4.0 International License

1. Introduction

Human preferences, farming practices, and agricultural environment all have an impact on the results of crop domestication, thereby decreasing mutations and variation. Selectivity driven by humans can be better understood due to developments in molecular technology involving genome sequencing [1]. Critics sometimes draw comparisons between conventional domestication and genetic modification (GM), but it's critical to recognize that these processes varies significantly in biological and social ways, which influences seed sovereignty and agrobiodiversity. Misrepresenting genetic modification as a continuation of domestication ignores the distinct mechanisms and consequences of each occurrence [2]. Through fixing nitrogen, boosting soil fertility, and enhancing water consumption efficiency, legumes benefit the environment and agricultural systems. Analyses on global food security and climate change are shifting from profitable species to underutilized ones with abundant genetic resources [3]. Crop evolution has been profoundly influenced by gene knockout alleles, from early agricultural origins to yield improvements during the Green Revolution. Recent genome sequencing has revealed advantageous loss-of-function occurrences in crop species, prompting efforts to accelerate crop breeding procedures by targeting gene knockouts with contemporary gene editing technologies [4]. Interest in de novo domestication of novel crops from wild species has increased with the latest advances in genome editing. De novo domestication must, be seen as an iterative process. Agronomic innovation, connections, and general genetic shifts as well as variations in human social preferences might all be significant [5]. Due to climatic change and geographic isolation, new species are formed, and genetic materials and accessions are essential for germplasm gathering. Competitor and predator effect diversification. Stress tolerance, yield, and quality are the main goals of plant breeding, that employs genetic diversity to enhance crops [6]. Genes related to seed qualities as dormancy, lifespan, germination, and vigor are identified by high-throughput gene expression analysis conducted on crops. Scientists have discovered unique genes that are active at various phases of seed formation. The analysis draws attention to the possible impact of epigenetic variables on seed growth and chemical composition, including imprinted genes in the seed-endosperm connection [7]. Faster gene discovery of agronomic factors influencing food production and quality has been made possible by computing resources, accessible genome sequencing, and molecular phenotyping technological advances [8]. For crop adaptation to climate change, crop intensification is essential, and genomics may accelerate the evolution of adaptable breeds. The acceptance of novel varieties and the effectiveness of resource-intensive breeding approaches may be impeded by centralized breeding initiatives' inability to satisfy the demands of smallholder farmers [9].

2. The Evolution of Crop Genetics

Though self-sufficiency is the objective of the food security strategy regarding seeds, it is dependent on imported seeds due to factors that include low quality, low competitiveness, insufficient infrastructure, antiquated machinery, and the absence of qualified professionals [10]. Biotechnology and genetic engineering have benefited significantly from genome editing and genetic selection approaches. For better commercial crops, shorten breeding time, promote novel genetic variations, and produce genetically modified plants with disease resistance attributes, molecular markers are employed in functional genomics, selection, editing, and gene targeting [11]. Although plant genetic resources (PGRs) are essential for agricultural resilience and profitability, population expansion is predicted to make them even more significant. However, due to genetic deterioration and environmental changes, numerous PGRs have disappeared, including landraces, wild relatives, and contemporary breeds [12]. Utilizing seed

retention in agricultural weed species including ryegrass, wild radish, and weedy amaranths, innovative weed management approaches like harvest weed seed control are implemented. By combining these approaches with genetics and molecular biology, seed shattering in weed and agricultural species may be minimized ^[13]. A greater knowledge of the interaction between genotype, phenotype, and environment is made possible by big data on crop characteristics, a massive dataset spanning many sizes and ecosystems. This improves plant molecular breeding, functional genomics, and effective farming techniques ^[14]. The global food security is dependent upon crop genetics. Since the *Arabidopsis thaliana* genome was sequenced, more than 100 crop genomes have been sequenced, and new technology and strategies might encourage more advancement in this field. Sequencing of the genomes of rice and *Arabidopsis thaliana* is a breakthrough ^[15]. Non-thermal plasma, also referred to as cold plasma, holds promise as an alternative to treat seeds including wheat, beans, corn, soybeans, barley, peanuts, rice, and *Arabidopsis thaliana* while also removing the necessity of pesticides and improving germination rates, crop shelf life, and ecological tolerance ^[16]. For plant populations to survive and grow, seed distribution is essential. Since the majority of plant species depend upon animals to distribute their seeds, modifications to external conditions that directly affect animals may have far-reaching consequences for plant ecosystems ^[17]. Physiological and demographic variables frequently impact plant adaptation to abiotic environments, but swift environmental changes may additionally affect seed dispersal processes. Considering ecological variety and phenotypic heterogeneity in characteristics associated to dispersal among seeds and their dispersers, monitoring these changes is complex ^[18]. As sea levels rise, the tidal marsh ecosystem is undergoing significant modifications. However, not much is known about the biological mechanisms driving this movement, especially with regard to the preservation and distribution of seeds in the soil ^[19]. Crop resilience, productivity, and adaptability are all endangered by crop genetic erosion, or the loss of crop variety. We still don't fully comprehend the scope, direction, reasons, and significance of these losses, despite warnings more than a century ago ^[20].

3. Recommendation

We propose the following recommendations for the future crop genomics.

- The ecology, physiology, and genetics of seed shattering should be the main topics of future exploration. Particular attention should be paid to the potential of weed species that are important to agriculture and the effects of environmental conditions.
- For agricultural production constraints to be addressed and future food security to be guaranteed, high-throughput genotypic and phenotypic approaches are essential for sustainable genetic resource exploitation.
- It is possible to improve future projections and estimate both the best- and worst-case scenarios by integrating theoretical and mathematical models with practical knowledge of seed dispersal mechanisms.
- Understanding the likelihood of species co-extinction and ecosystem buffering is possible through observational research and rewiring models. In order to address de-faunation impacts, it is imperative to comprehend methods for compensating for seed disperser loss.
- To combat climate change and population expansion, crop development is essential. Both improved genotype selection and genetic variety are necessary for success. In traditional crosses, animals that have close connections with one another could develop greater genetic variety, and induced mutations can provide unique genetic traits.


- Similar to gene editing and crucial decision-making in domestication pathways, new food crop domestication is a complex public health endeavor. Even if vaccinations are appealing, advancements in vaccination acceptability must include ongoing surveillance of viral development and transmission, improvements in vaccine delivery technology, support for clinics and vaccination campaigns, and cultural outreach.
- Technological developments in genetics and molecular biology have the ability to manage weed species by decreasing agricultural seed shattering. Further research need to investigate various processes of seed splitting in connection with climate and farming methods.

Conclusion

Through effective management techniques and the maintenance of unique seed trait variants within gene banks, genetic and genomic resources facilitate the development of precise cultivars and offer important insights into the biology and attributes of seeds. When assessing the productivity and financial advantages of food crops, seed production is an essential indicator. Breeding efforts are directed on minimizing seed cracking, a biological trait present in many weed species. In many crop kinds, seed breaking is still a major problem even if human interventions have decreased it in food crops. The function of gene LoF in rapid evolution and adaptability—particularly in domestication and crop species augmentation—is being investigated by evolutionary biologists. This leads to the development of gene editing techniques for particular crop growth. The “Second Green Revolution” of modern agriculture is under threat because the major yield increases of the Green Revolution were mostly brought about by spontaneous LoF mutations. On seed germination, plant development, yield, and stress tolerance, researchers are investigating the effects of non-thermal plasma (NTP). Water contact angle, DNA methylation, gene expression patterns, germination speed, seedling development, enzyme activity, hormone presence, and the transmission of enhanced features to subsequent generations are just a few of the impacts of NTP on seeds that are being explored.

ORCID iDs

Mansi Sahu  <https://orcid.org/0009-0003-2859-0620>

Kailash Bisht  <https://orcid.org/0000-0003-3659-2012>

Sanjeev Kumar Shah  <https://orcid.org/0000-0002-9978-5842>

References

1. Smýkal P., Nelson M. N., Berger J. D., Von Wettberg E. J. (2018) The impact of genetic changes during crop domestication *Agronomy* 7(8)119.
2. Mueller N. G., Flachs A. (2022) Domestication, crop breeding, and genetic modification are fundamentally different processes: implications for seed sovereignty and agrobiodiversity *Agriculture and Human Values* 1(39)455–472.
3. Sahrzaini N. A., Rejab N. A., Harikrishna J. A., Khairul Ikram N. K., Ismail I., Kugan H. M. (2020) Pulse crop genetics for a sustainable future: Where we are now and where we should be heading *Frontiers in plant science* 11532119.
4. Monroe J. G., Arciniegas J. P., Moreno J. L., Sánchez F., Sierra S., Valdes S... Chavarriaga P. (2020) The lowest hanging fruit: Beneficial gene knockouts in past, present, and future crop evolution *Current Plant Biology* 24100185.
5. Van Tassel D. L., Tesdell O., Schlautman B., Rubin M. J., DeHaan L. R., Crews T. E., Streit Krug A. (2020) New food crop domestication in the age of gene editing: genetic, agronomic and cultural change remain co-evolutionarily entangled *Frontiers in plant science* 11524819.

6. Begna T., Begna T. (2021) Role and economic importance of crop genetic diversity in food security *International Journal of Agricultural Science and Food Technology* 1(7)164–169.
7. Dwivedi S. L., Spillane C., Lopez F., Ayele B. T., Ortiz R. (2021) First the seed: Genomic advances in seed science for improved crop productivity and food security *Crop Science* 3(61)1501–1526.
8. Steinwand M. A., Ronald P. C. (2020) Crop biotechnology and the future of food *Nature Food* 5(1)273–283.
9. Fadda C., Mengistu D. K., Kidane Y. G., Dell'Acqua M., Pè M. E., Van Etten J. (2020) Integrating conventional and participatory crop improvement for smallholder agriculture using the seeds for needs approach: A review *Frontiers in Plant Science* 11559515.
10. Zolkin A. L., Matvienko E. V., Shavanov M. V. (2021, March) Innovative technologies in agricultural crops breeding and seed farming In *IOP Conference Series: Earth and Environmental Science* (Vol.No. 2,022092 IOP Publishing 677 p.
11. Dheer P., Rautela I., Sharma V., Dhiman M., Sharma A., Sharma N., Sharma M. D. (2020) Evolution in crop improvement approaches and future prospects of molecular markers to CRISPR/Cas9 system *Gene* 753144795.
12. Salgotra R. K., Chauhan B. S. (2023) Genetic diversity, conservation, and utilization of plant genetic resources *Genes* 1(14)174.
13. Maity A., Lamichaney A., Joshi D. C., Bajwa A., Subramanian N., Walsh M., Bagavathiannan M. (2021) Seed shattering: a trait of evolutionary importance in plants *Frontiers in Plant Science* 12657773.
14. Fan J., Zhang Y., Wen W., Gu S., Lu X., Guo X. (2021) The future of Internet of Things in agriculture: Plant high-throughput phenotypic platform *Journal of Cleaner Production* 280123651.
15. Purugganan M. D., Jackson S. A. (2021) Advancing crop genomics from lab to field *Nature genetics* 5(53)595–601.
16. Starič P., Vogel-Mikuš K., Mozetič M., Junkar I. (2020) Effects of nonthermal plasma on morphology, genetics and physiology of seeds: A review *Plants* 12(9)1736.
17. Rogers H. S., Donoso I., Traveset A., Fricke E. C. (2021) Cascading impacts of seed disperser loss on plant communities and ecosystems *Annual Review of Ecology, Evolution, and Systematics* 52641–666.
18. Johnson J. S., Cantrell R. S., Cosner C., Hartig F., Hastings A., Rogers H. S.... Pufal G. (2019) Rapid changes in seed dispersal traits may modify plant responses to global change *AoB Plants* 3(11)plz020.
19. Kottler E. J., Gedan K. (2020) Seeds of change: characterizing the soil seed bank of a migrating salt marsh *Annals of Botany* 2(125)335–344.
20. Khoury C. K., Brush S., Costich D. E., Curry H. A., De Haan S., Engels J. M.... Thormann I. (2022) Crop genetic erosion: understanding and responding to loss of crop diversity *New Phytologist* 1(233)84–118.